

University of Groningen

Water abundances in high-mass protostellar envelopes

Marseille, M. G.; van der Tak, F. F. S.; Herpin, F.; Wyrowski, F.; Chavarria, L.; Pietropaoli, B.; Baudry, A.; Bontemps, S.; Cernicharo, J.; Jacq, T.

Published in:
Astronomy & astrophysics

DOI:
[10.1051/0004-6361/201015103](https://doi.org/10.1051/0004-6361/201015103)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Marseille, M. G., van der Tak, F. F. S., Herpin, F., Wyrowski, F., Chavarria, L., Pietropaoli, B., Baudry, A., Bontemps, S., Cernicharo, J., Jacq, T., Frieswijk, W., Shipman, R., van Dishoeck, E. F., Bachiller, R., Benedettini, M., Benz, A. O., Bergin, E., Bjerkeli, P., Blake, G. A., ... Yildiz, U. A. (2010). Water abundances in high-mass protostellar envelopes: Herschel observations with HIFI. *Astronomy & astrophysics*, 521, [L32]. <https://doi.org/10.1051/0004-6361/201015103>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/47376038>

Water abundances in high-mass protostellar envelopes: Herschel observations with HIFI

Article in *Astronomy and Astrophysics* · July 2010

DOI: 10.1051/0004-6361/201015103 · Source: OAI

CITATIONS

12

READS

41

9 authors, including:



Fabrice Herpin

Université Bordeaux 1

129 PUBLICATIONS 2,161 CITATIONS

[SEE PROFILE](#)



Lourdes María Chavarria

University of São Paulo

44 PUBLICATIONS 341 CITATIONS

[SEE PROFILE](#)



Bastien Pietropaoli

University College Cork

20 PUBLICATIONS 99 CITATIONS

[SEE PROFILE](#)



A. Baudry

University of Bordeaux

278 PUBLICATIONS 2,236 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



ATLASGAL [View project](#)



Magnetic activity in the extended photospheres of AGB stars [View project](#)

Water abundances in high-mass protostellar envelopes: Herschel observations with HIFI[★]

M.G. Marseille¹⁰, F.F.S. van der Tak^{10,11}, F. Herpin⁶, F. Wyrowski³⁰, L. Chavarría⁶, B. Pietropaoli⁶, A. Baudry⁶, S. Bontemps⁶, J. Cernicharo¹⁶, T. Jacq⁵, W. Frieswijk¹⁰, R. Shipman¹⁰, E.F. van Dishoeck^{1,2}, R. Bachiller¹², M. Benedettini¹³, A.O. Benz³, E. Bergin¹⁴, P. Bjerkeli⁹, G.A. Blake¹⁵, J. Braine⁶, S. Bruderer³, P. Caselli^{4,5}, E. Caux^{47,48}, C. Codella⁵, F. Daniel¹⁶, P. Dieleman¹⁰, A.M. di Giorgio¹³, C. Dominik^{17,18}, S.D. Doty¹⁹, P. Encrenaz²⁰, M. Fich²¹, A. Fuente²², T. Gaier³¹, T. Giannini²³, J.R. Goicoechea¹⁶, Th. de Graauw¹⁰, F. Helmich¹⁰, G.J. Herczeg², M.R. Hogerheijde¹, B. Jackson¹⁰, H. Javadi³¹, W. Jellema¹⁰, D. Johnstone^{7,8}, J.K. Jørgensen²⁴, D. Kester¹⁰, L.E. Kristensen¹, B. Larsson²⁵, W. Laauwen¹⁰, D. Lis²⁶, R. Liseau⁹, W. Luinge¹⁰, C. McCoey^{21,27}, A. Megej⁴⁶, G. Melnick²⁸, D. Neufeld²⁹, B. Nisini²³, M. Olberg⁹, B. Parise³⁰, J.C. Pearson³¹, R. Plume³², C. Risacher¹⁰, P. Roelfsema¹⁰, J. Santiago-García³³, P. Saraceno¹³, P. Siegel³¹, J. Stutzki⁴³, M. Tafalla¹², T.A. van Kempen²⁸, R. Visser¹, S.F. Wampfler³, and U.A. Yıldız¹

(Affiliations can be found after the references)

Received xxx; accepted xxx

ABSTRACT

Aims. We derive the **dense core structure and the water abundance** in four massive star-forming regions **which may help understand the earliest stages of massive star formation.**

Methods. We present Herschel-HIFI observations of the para-H₂O 1₁₁ – 0₀₀ and 2₀₂ – 1₁₁ and the para-H₂¹⁸O 1₁₁ – 0₀₀ transitions. The envelope contribution to the line profiles is separated from contributions by outflows and foreground clouds. The envelope contribution is modeled using Monte-Carlo radiative transfer codes for dust and molecular lines (MC3D and RATRAN), with the water abundance and the turbulent velocity width as free parameters.

Results. While the outflows are mostly seen in emission in high-J lines, **envelopes are seen in absorption in ground-state lines**, which are almost saturated. The derived water abundances range from 5×10^{-10} to 4×10^{-8} **in the outer envelopes. We detect cold clouds surrounding the protostar envelope, thanks to the very high quality of the Herschel-HIFI data and the unique ability of water to probe them.** Several foreground clouds are also detected along the line of sight.

Conclusions. The low H₂O abundances in massive dense cores are in accordance with the expectation that high densities and low temperatures lead to freeze-out of water on dust grains. **The spread in abundance values is not clearly linked to physical properties of the sources.**

Key words. ISM: dust, extinction – ISM: molecules – ISM: abundances

1. Introduction

Massive stars ($\gtrsim 10 M_{\odot}$) play a major role in the interstellar energy budget and the shaping of the Galactic environment (Zinnecker & Yorke 2007). **However, the formation of such high-mass stars is not well understood due to several reasons: they are rare, have a short evolution time scale, they are born deeply embedded, and are far from the solar system.**

The main sequence lifetime of massive stars is preceded by an embedded phase which subdivides into several classes of objects: massive pre-stellar cores (mPSC), which are local temperature minima and density maxima within dark clouds (Sridharan et al. 2005); high-mass protostellar objects (HMPO), where a central protostar is surrounded by a massive envelope with a centrally peaked temperature and density distribution (van der Tak et al. 2000); hot molecular cores (HMC), which have larger masses of warm gas and dust, and high abun-

dances of complex organic molecules which have evaporated off dust grains and/or formed by warm gas-phase chemistry (Motte et al. 2003); and ultracompact HII regions (UCHII), which show large pockets of ionized gas confined to the star (Churchwell et al. 1990). A key question is to what extent these phases represent differences in luminosity and/or age, and if all high-mass stars pass through all these phases.

The water molecule is thought to be a sensitive tracer of physical conditions in star-forming regions, which acts as a natural filter for warm gas **because of its large abundance variations between hot and cold regions** (van der Tak et al. 2006). **Moreover, because the dust continuum is strong at the higher frequencies, water lines connecting with the lowest energy levels can be seen in absorption, thus providing an alternative method to probe different depths in the protostellar environment** (Poelman & van der Tak 2007). Measurements of the abundance of water are therefore a step toward understanding the energy budget of star-forming regions, and thus of the star formation process itself.

[★] Herschel is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with participation of NASA.

This paper presents water observations performed with the Heterodyne Instrument for the Far-Infrared (HIFI; de Graauw & et al. 2010) on-board ESA's Herschel Space Observatory (Pilbratt & et al. 2010). We use the p-H₂O ground-state line and two lines which constrain the excitation and optical depth (Table 1), **all three lying at similar frequencies and observed at similar resolution**. The sources are four massive star-forming regions (Table 2): the HMCs G31.41+0.31 and G29.96−0.02 and the HMPOs W33A and W43-MM1. We compare our results with those for two other regions: the UCHII region DR21 (van der Tak et al. 2010) and the HMPO W3 IRS5 (Chavarría et al. 2010, this volume).

However, the aim is to discover trends in the water line emission for future extended studies, identifying links in the water abundance between the various evolutionary stages of high-mass star formation and to use water as probe of the gas dynamics around protostars. Given the small number of sources and lines observed, it is premature to look for general trends. The large amount of upcoming Herschel-HIFI data will help on this issue.

Table 1. List of lines.

Molecule	Transition	ν (GHz)	E_{up} (K)	n_{crit}^a (cm ^{−3})	σ_{rms} (mK)
H ₂ O	1 ₁₁ − 0 ₀₀	1113.343	53.4	1.7×10^8	40
H ₂ O	2 ₀₂ − 1 ₁₁	987.927	100.8	2.1×10^8	50
H ₂ ¹⁸ O	1 ₁₁ − 0 ₀₀	1101.698	53.4	1.7×10^8	40

Notes. ^(a) Values at 20 K from collision rates of Grosjean et al. (2003)

Table 2. List of sources.

Name	R.A. J2000	Dec. J2000	L (10 ⁴ L _⊙)	d^a (kpc)	V_{LSR} (km s ^{−1})
G31.41+0.31	18 ^h 47 ^m 34.3 ^s	−01°12′46.0″	15	7.9	+98.8
G29.96−0.02	18 ^h 46 ^m 03.8 ^s	−02°39′22.0″	20	7.4	+98.7
W33A	18 ^h 14 ^m 39.1 ^s	−17°52′07.0″	8.5	4.0	+37.5
W43-MM1	18 ^h 47 ^m 47.0 ^s	−01°54′28.0″	2.2	5.5	+98.8

Notes. ^(a) Values from Hatchell & van der Tak (2003), except W43-MM1 (Motte et al. 2003) and W33A (van der Tak et al. 2000).

2. Observations

The four regions have been observed with HIFI on the 3rd, 4th and 6th of March 2010 (see Table 2). Spectra were taken in double sideband mode using receivers 4a (p-H₂O at 988 GHz) and 4b (p-H₂O and p-H₂¹⁸O at 1113 GHz and 1102 GHz) with ν_{LO} = 980 GHz and 1108 GHz respectively. The observations are part of the Priority Science Program (PSP) of the Guaranteed-Time Key Program *Water In Star-forming regions with Herschel* (WISH; Van Dishoeck et al., in prep.).

Data were simultaneously taken with the acousto-optical Wide-Band Spectrometer (WBS) and the correlator-based High-Resolution Spectrometer (HRS), in both horizontal and vertical polarizations. **This paper focuses on data from the WBS**, which covers 1140 MHz bandwidth at 1.1 MHz spectral resolution (~ 0.3 km s^{−1}) (Roelfsema et al. 2010). System temperatures range between 350 K around 1113 GHz and 450 K

around 988 GHz; receiver 4a in V polarization shows particularly high values. Integration times (ON+OFF) were 193 s for the 1113 GHz and the 1102 GHz lines, and 206 s for the 988 GHz line, for each source, and the rms noise levels reached are 40–50 mK (Table 1). Observations have been reduced with the Herschel Interactive Processing Environment¹ (HIPE) version 2.8. The intensity scale is converted to T_{mb} using main beam efficiencies of 0.74. The double-side band continuum level has been divided by 2 to make its brightness directly comparable to that of the lines, which are measured in single sideband.

3. Results

Observed water lines for the four studied regions are shown in Fig. 1. The H₂O 1₁₁ − 0₀₀ line shows an absorption at the systemic velocity (V_{LSR}) in all sources. In all cases except G31.41+0.31, outflow wings are detected close to the main absorption, with a maximal shift of 3 km s^{−1}. These wings are seen in emission, which indicates an origin in hot, low-density (10^3 cm^{−3}) gas (Poelman & van der Tak 2007). **Absorption features are seen over a large velocity range in G29.96−0.02, W33A, W43-MM1 and more weakly in G31.41+0.31.** The absorptions at velocity offsets >4 km s^{−1} likely originate in cold foreground clouds on the line of sight to the source. In contrast, the absorption features at smaller velocity offsets **are plausibly related** to cold clouds surrounding the dense cores (which other studies call the protostellar envelopes), which are all part of large-scale molecular clouds (see Fig. 2). **Table 3 presents a Gaussian decomposition of the line profiles around V_{LSR} ; Appendix A shows Gaussian iterative decompositions of the absorption profiles of the ground-state transition over the full velocity range, showing a large number of velocity components thanks to the high velocity resolution of the HIFI instrument.** The absorptions at V_{LSR} are saturated for G31.41+0.31 and W33A and nearly saturated for the other sources, which indicates abundances around $\sim 10^{-9}$ for the outer cold parts of the massive dense cores (Poelman & van der Tak 2007).

The H₂O 2₀₂ − 1₁₁ line always appears in emission and shows a broad and a narrower velocity component (Fig. 1). In addition, the spectra of G31.41+0.31 and W43-MM1 show two well defined self-absorption features which appear at the source velocity. With its high E_{up} , this transition mainly traces warm gas, and the presence of these absorption features in G31.41+0.31 and W43-MM1 suggests a higher water abundance in these sources than in G29.96−0.02 and W33A. The components seen in emission have Gaussian shapes, one being wider (FWHM=20–40 km s^{−1}) than the other (FWHM=6.4–8.0 km s^{−1}). We associate the broad component with high-velocity outflows **associated with the protostar also seen in 1₁₁ − 0₀₀ line emission**. This component is symmetric with respect to the source velocity in G29.96−0.02 and W43-MM1, blueshifted by 2.8 km s^{−1} in W33A, and redshifted by 4.4 km s^{−1} in G31.41+0.31 (Table 3). The narrower (hereafter ‘medium’) component **is potentially** associated with shocked surrounding material where water is released in the gas phase. Indeed, shocks occur at the interface between jets and the surrounding dense envelope, with a velocity close to that of the massive dense core. Similar results are found in Kristensen et al. (2010), Johnstone et al. (2010) and Chavarría et al. (2010).

The H₂¹⁸O 1₁₁ − 0₀₀ transition is seen in absorption at the source velocity in G31.41 and W43-MM1, which is not sat-

¹ <http://herchel.esac.esa.int/>




Fig. 1. Herschel/HIFI spectra of the $\text{H}_2\text{O } 1_{11} - 0_{00}$ (top), $\text{H}_2\text{O } 2_{02} - 1_{11}$ (middle) and $\text{H}_2^{18}\text{O } 1_{11} - 0_{00}$ (bottom) lines. Dashed lines drawn at V_{LSR} .

Table 3. Gaussian decomposition of the line profiles at velocities close to V_{LSR} .

Source	para- $\text{H}_2\text{O } (1_{11} - 0_{00})$			para- $\text{H}_2\text{O } (2_{02} - 1_{11})$			para- $\text{H}_2^{18}\text{O } (1_{11} - 0_{00})$		
	V_{LSR} (km s^{-1})	T_{mb} (K)	ΔV (km s^{-1})	V_{LSR} (km s^{-1})	T_{mb} (K)	ΔV (km s^{-1})	V_{LSR} (km s^{-1})	T_{mb} (mK)	ΔV (km s^{-1})
G31.41+0.31	95.1	0.94*	3.7	94.6	1.37	6.4	99.5	0.27*	5.2
				99.3	0.42*	14.0			
				103.7	0.2	~40			
G29.96-0.02	91.3	0.26*	3.9	97.8	1.10	21	98.5	290	6.0
	98.5	0.90	18.8	98.2	3.21	8.0			
	99.4	0.99*	8.4						
	103.2	0.40*	2.3						
W33A	35.9	0.85*	11.0	35.5	0.34	28	34.2	0.23*	11.8
	43.0	0.53	20.0	38.3	1.87	7.0	37.3	280	8.4
W43-MM1	98.7	0.87*	13.5	99.6	0.89	22	99.4	0.19*	8.7
	103.3	0.43	14.0	99.7	0.31*	6.8			

Notes. (*) Absorption lines are indicated in $T_{\text{abs}}/T_{\text{continuum}}$ scales.

urated. This feature originates from the massive envelope. In G29.96, the pure emission profile of this transition implies a warm diffuse gas origin. Since G29.96 is also the brightest source in the narrower component of the 988 GHz line, we suggest that the H_2^{18}O emission is dominated by shocks at the interface between jets and the envelope. The P-Cygni-like profile for W33A is a mix of both behaviors: the sum of an absorption feature due to the massive core and emission from an outflow. The fact that the emission is seen only on the red-shifted part of the profile is consistent with the outflow components seen in 1113 GHz and 988 GHz lines, which are also more powerful on the red part of the spectrum.

4. Discussion and conclusions

To derive the water abundance in the four massive dense cores we have removed features related to outflows and foreground clouds from the spectrum before any line modeling. The high spectral resolution of HIFI is essential in this process, in particular for the absorbers with velocities close to that of the central source. Studying the $\text{H}_2^{18}\text{O } 1_{11} - 0_{00}$ transition prior to the others also facilitates to disentangle the **envelope** contribution, **since this line is not saturated having a lower optical depth than the main H_2O isotope.**

Once the main contribution is extracted, we model its profile according to the method described in Marseille et al. (2008): first, the dust emission from the massive dense core is reproduced with the MC3D radiative transfer code (Wolf et al. 1999), including total luminosity and density profile from the literature (power-law index $p = -1.5$); second, the temperature profile obtained is used to model the line emission with the RATRAN code (Hogerheijde & van der Tak 2000). The free parameters are: $X_{\text{H}_2\text{O}}$ the molecular abundance relative to H_2 , and v_{turb} , the turbulent velocity width.

Good fits are obtained for the $\text{H}_2^{18}\text{O } 1_{11}-0_{00}$ transition, which is not saturated unlike the H_2O lines. The fitting considers both the line strength (area and width) and the profile shapes. We have computed a grid of $X_{\text{H}_2\text{O}}$ and v_{turb} values, adapting step by step the grid around the best χ^2 . Using a $^{16}\text{O}/^{18}\text{O}$ ratio of 500, we proceed to model the main isotopic water lines. The H_2O abundance is kept constant in our models. We have tried models with an abundance increase in the inner region where $T > 100$ K, but the current data do not favor these models above the constant-abundance models.

We estimate the absolute uncertainty in the retrieved H_2O abundance to be a factor of 10. Since we use the same modeling strategy as the studies by van der Tak et al. (2010) and Chavarría et al. (2010), the **abundances obtained** should be comparable to **better than a factor of 3**. Our observed spread in abundances of a factor of ~ 100 is much larger than this uncertainty. The same range of abundances is found in other HIFI-based studies of high-mass star-forming regions (van der Tak et al. 2010; Chavarría et al. 2010), and also in previous work with ISO (Boonman et al. 2003) and from the ground (van der Tak et al. 2006).

In conclusion, for the massive star forming regions described in this letter, we clearly detect the contribution of the envelope within the dense core. It is limited to a strong self-absorbed feature, mainly seen in the ground-state line. In order to evaluate it, we first have to remove emission from outflow shocks and absorption by foreground clouds along the line of sight. The velocities of the absorbers indicate that some are part of the close environment of the source, while others are physically unrelated. The derived massive dense core abundances suggest a strong freeze-out of water on dust grains, and imply that water plays only a minor role in the thermal balance of the gas.

The H_2O line profiles do not seem to depend on the supposed evolutionary stage of the source. For example, the two ‘hot molecular cores’ G31.41 and G29.96 show very different line profiles, and also their H_2O abundances differ by a factor of ~ 100 . Also, the abundance variations that we have found do not seem related to the luminosity of the sources, their temperature or their turbulent velocity field. However, the number of cases treated is not sufficient for a statistical treatment. Future studies following the same procedure with a larger number of sources should conclude on this issue. Within our sample, the highest H_2O abundances are derived for G31.41 and W43-MM1 which show self-absorbed $2_{02}-1_{11}$ line profiles (Fig. 1). As these sources are not the most luminous, hot, or active ones in our sample, the origin of such a high abundance is unclear.

Firm conclusions about a link between water emission behaviour and the evolutionary stage of the source are limited by the small number of sources. Our data show that water is a useful tool to understand the gas dynamics in and around massive stars forming regions. Future multi-line studies of larger samples are highly promising to answer key questions about the formation of massive stars.

Acknowledgements. HIFI has been designed and built by a consortium of institutes and university departments from across Europe, Canada and the United States under the leadership of SRON Netherlands Institute for Space Research, Groningen, The Netherlands and with major contributions from Germany, France and the US. Consortium members are: Canada: CSA, U.Waterloo; France: CESR, LAB, LERMA, IRAM; Germany: KOSMA, MPIfR, MPS; Ireland, NUI Maynooth; Italy: ASI, IFSI-INAF, Osservatorio Astrofisico di Arcetri- INAF; Netherlands: SRON, TUD; Poland: CAMK, CBK; Spain: Observatorio Astronómico Nacional (IGN), Centro de Astrobiología (CSIC-INTA). Sweden: Chalmers University of Technology - MC2, RSS & GARD; Onsala Space Observatory; Swedish National Space Board, Stockholm University - Stockholm Observatory; Switzerland: ETH Zurich, FHNW; USA: Caltech, JPL, NHSC.

HIFE is a joint development by the Herschel Science Ground Segment Consortium, consisting of ESA, the NASA Herschel Science Center, and the HIFI, PACS and SPIRE consortia.

References

- Boonman, A., Doty, S., van Dishoeck, E., et al. 2003, A&A, 406, 937
- Chavarría, L., Herpin, F., Jacq, T., & the WISH consortium. 2010, A&A, in press
- Churchwell, E., Walmsley, C. M., & Cesaroni, R. 1990, A&As, 83, 119
- de Graauw & et al. 2010, A&A, in press
- Grosjean, A., Dubernet, M., & Ceccarelli, C. 2003, A&A, 408, 1197
- Hatchell, J. & van der Tak, F. F. S. 2003, A&A, 409, 589
- Hogerheijde, M. R. & van der Tak, F. F. S. 2000, A&A, 362, 697
- Johnstone, D., Fish, M., M^CCoey, C., et al. 2010, A&A, in press
- Kristensen, L., Visser, R., van Dishoeck, E., et al. 2010, A&A, in press
- Marseille, M., Bontemps, S., Herpin, F., et al. 2008, A&A, 488, 579
- Motte, F., Schilke, P., & Lis, D. C. 2003, ApJ, 582, 277
- Pilbratt & et al. 2010, A&A, in press
- Poelman, D. R. & van der Tak, F. F. S. 2007, A&A, 475, 949
- Roelfsema, P., Helmich, F., Teyssier, D., et al. 2010, A&A, in press
- Sridharan, T. K., Beuther, H., Saito, M., Wyrowski, F., & Schilke, P. 2005, ApJ, 634, L57
- van der Tak, F., Marseille, M., Herpin, F., et al. 2010, A&A, in press
- van der Tak, F., van Dishoeck, E., Evans, I. N., & Blake, G. 2000, ApJ, 537, 283
- van der Tak, F., Walmsley, C., Herpin, F., et al. 2006, A&A, 447, 1011
- Wolf, S., Henning, T., & Stecklum, B. 1999, A&A, 349, 839
- Zinnecker, H. & Yorke, H. W. 2007, ARA&A, 45, 481




Fig. 2. Extraction of the saturated absorption of para-H₂O 1₁₁–0₀₀ line in W43-MM1. Original profile appears in black bold, residual in green bold.

Table 4. Model parameters and derived water abundances.

Source	M_{gas} (M_{\odot})	r_{min} (AU)	r_{max} (AU)	$n(r_{\text{min}})$ (cm^{-3})	$n(r_{\text{max}})$ (cm^{-3})	$T(r_{\text{min}})$ (K)	$T(r_{\text{max}})$ (K)	$X_{\text{H}_2\text{O}}$	v_{turb} (km s^{-1})
G31.41+0.31	1500	200	22 515	8.1×10^8	3.1×10^6	406	43.2	3.1×10^{-8}	1.4
G29.96-0.02	700	200	20 700	4.4×10^8	1.9×10^6	489	50.8	$< 5.0 \times 10^{-10}$	1.1
W33A	4000	200	62 000	3.5×10^8	4.0×10^5	291	26.0	6.0×10^{-10}	1.6
W43-MM1	2000	200	27 500	5.0×10^8	2.3×10^6	243	24.7	4.0×10^{-8}	3.0
DR21 ^a	1650	2000	60 520	1.6×10^7	1.5×10^5	117	23.3	2.0×10^{-10}	3.0
W3-IRS5 ^b	250	200	12 000	2.9×10^8	2.7×10^6	480	54.7	2.0×10^{-8}	2.0

Notes. ^(a) Values from van der Tak et al. (2010) ^(b) Values from Chavarría et al. (2010)

Appendix A: Massive dense core component extraction

The velocity profiles of the H₂O 1₁₁–0₀₀ line show absorption features at several velocities. These absorption features arise in foreground clouds along the line of sight or in cold clouds in the neighbourhood of the massive dense core, and are not saturated unlike the absorption from the massive **envelope**. In addition to these absorptions, some sources show H₂O emission from protostellar outflows.

This appendix presents our procedure to remove these features in order to extract the contribution from the envelope to the line profile. Contrary to others, **absorption** from this part of the object is saturated. We are then able to remove other features by iterative Gaussian fits. This process is helped by the high velocity resolution provided by the Herschel-HIFI instrument, showing accurate and "bumpy" profiles in absorptions. Assuming that each bump corresponds to a velocity component, they are removed using the Gaussian fitting tool available in the HIPE software. Starting from the component with the lowest velocity, they are extracted one by one, using the residual of the previous removal to fit the next one. This way of fitting insures a very good extraction of velocity component, giving a quasi-unique final decomposition of the absorption features. Results of this process are given in Fig. 2, A.1, A.2, A.3 and Tables ??, ??, ?? and ??.

- ¹ Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands
- ² Max Planck Institut für Extraterrestrische Physik, Giessenbachstrasse 1, 85748 Garching, Germany
- ³ Institute of Astronomy, ETH Zurich, 8093 Zurich, Switzerland
- ⁴ School of Physics and Astronomy, University of Leeds, Leeds LS2 9JT, UK
- ⁵ INAF - Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125 Firenze, Italy
- ⁶ Université de Bordeaux, Laboratoire d'Astrophysique de Bordeaux, France; CNRS/INSU, UMR 5804, Floirac, France
- ⁷ National Research Council Canada, Herzberg Institute of Astrophysics, 5071 West Saanich Road, Victoria, BC V9E 2E7, Canada
- ⁸ Department of Physics and Astronomy, University of Victoria, Victoria, BC V8P 1A1, Canada
- ⁹ Department of Radio and Space Science, Chalmers University of Technology, Onsala Space Observatory, 439 92 Onsala, Sweden
- ¹⁰ SRON Netherlands Institute for Space Research, PO Box 800, 9700 AV, Groningen, The Netherlands
- ¹¹ Kapteyn Astronomical Institute, University of Groningen, PO Box 800, 9700 AV, Groningen, The Netherlands
- ¹² Observatorio Astronómico Nacional (IGN), Calle Alfonso XII, 3. 28014, Madrid, Spain
- ¹³ INAF - Istituto di Fisica dello Spazio Interplanetario, Area di Ricerca di Tor Vergata, via Fosso del Cavaliere 100, 00133 Roma, Italy
- ¹⁴ Department of Astronomy, The University of Michigan, 500 Church Street, Ann Arbor, MI 48109-1042, USA
- ¹⁵ California Institute of Technology, Division of Geological and Planetary Sciences, MS 150-21, Pasadena, CA 91125, USA
- ¹⁶ Centro de Astrobiología. Departamento de Astrofísica. CSIC-INTA. Carretera de Ajalvir, Km 4, Torrejón de Ardoz. 28850, Madrid,




Fig. A.1. Extraction of the saturated absorption of para-H₂O $1_{11} - 0_{00}$ line in W33A. Original profile appears in black bold, residual in green bold.




Fig. A.2. Extraction of the saturated absorption of para-H₂O $1_{11} - 0_{00}$ line in G29.96. Original profile appears in black bold, residual in green bold.






Fig. A.3. Extraction of the saturated absorption of para-H₂O $1_{11} - 0_{00}$ line in G31.41. Original profile appears in black bold, residual in green bold.


G29.96-0.02

WBS


HRS




G29.96-0.02



G29.96-0.02





G31.41+0.31




G31.41+0.31

WBS

HRS





W33A




W33A

WBS

HRS



W43-MM1



W43-MM1

WBS

HRS

